


Lesson 1: The Pythagorean Theorem

Classwork


Example 1

Write an equation that will allow you to determine the length of the unknown side of the right triangle.




Example 2

Write an equation that will allow you to determine the length of the unknown side of the right triangle.




Example 3

Write an equation to determine the length of the unknown side of the right triangle.




Example 4


In the figure below, we have an equilateral triangle with a height of 10 inches. What do we know about an equilateral triangle?

**Exercises 1–3**


1. Use the Pythagorean Theorem to estimate the length of the unknown side of the right triangle. Explain why your estimate makes sense.



2. Use the Pythagorean Theorem to estimate the length of the unknown side of the right triangle. Explain why your estimate makes sense.



3. Use the Pythagorean Theorem to estimate the length of the unknown side of the right triangle. Explain why your estimate makes sense.



Lesson Summary

Perfect square numbers are those that are a product of an integer factor multiplied by itself. For example, the number 25 is a perfect square number because it is the product of 5 multiplied by 5.

When the square of the length of an unknown side of a right triangle is not equal to a perfect square, you can estimate the length by determining which two perfect squares the number is between.

Example:



Let c represent the length of the hypotenuse. Then,

$$3^2 + 7^2 = c^2$$


$$9 + 49 = c^2$$

$$58 = c^2$$


The number 58 is not a perfect square, but it is between the perfect squares 49 and 64. Therefore, the length of the hypotenuse is between 7 and 8, but closer to 8 because 58 is closer to the perfect square 64 than it is to the perfect square 49.

Problem Set


- Use the Pythagorean Theorem to estimate the length of the unknown side of the right triangle. Explain why your estimate makes sense.




2. Use the Pythagorean Theorem to estimate the length of the unknown side of the right triangle. Explain why your estimate makes sense.




3. Use the Pythagorean Theorem to estimate the length of the unknown side of the right triangle. Explain why your estimate makes sense.




4. Use the Pythagorean Theorem to estimate the length of the unknown side of the right triangle. Explain why your estimate makes sense.




5. Use the Pythagorean Theorem to estimate the length of the unknown side of the right triangle. Explain why your estimate makes sense.




6. Determine the length of the unknown side of the right triangle. Explain how you know your answer is correct.




7. Use the Pythagorean Theorem to estimate the length of the unknown side of the right triangle. Explain why your estimate makes sense.



8. The triangle below is an isosceles triangle. Use what you know about the Pythagorean Theorem to determine the approximate length of base of the isosceles triangle.



9. Give an estimate for the area of the triangle shown below. Explain why it is a good estimate.




Lesson 2: Square Roots

Classwork

Exercises 1–4

1. Determine the positive square root of 81, if it exists. Explain.
2. Determine the positive square root of 225, if it exists. Explain.
3. Determine the positive square root of -36 , if it exists. Explain.
4. Determine the positive square root of 49, if it exists. Explain.

Discussion



Exercises 5–9

Determine the positive square root of the number given. If the number is not a perfect square, determine which integer the square root would be closest to, then use “guess and check” to give an approximate answer to one or two decimal places.

5. $\sqrt{49}$

6. $\sqrt{62}$

7. $\sqrt{122}$

8. $\sqrt{400}$

9. Which of the numbers in Exercises 5–8 are not perfect squares? Explain.

Lesson Summary

A positive number whose square is equal to a positive number b is denoted by the symbol \sqrt{b} . The symbol \sqrt{b} automatically denotes a positive number. For example, $\sqrt{4}$ is always 2, not -2 . The number \sqrt{b} is called a positive square root of b .

Perfect squares have square roots that are equal to integers. However, there are many numbers that are not perfect squares.

Problem Set

Determine the positive square root of the number given. If the number is not a perfect square, determine the integer to which the square root would be closest.

- $\sqrt{169}$
- $\sqrt{256}$
- $\sqrt{81}$
- $\sqrt{147}$
- $\sqrt{8}$
- Which of the numbers in Problems 1–5 are not perfect squares? Explain.
- Place the following list of numbers in their approximate locations on a number line:

$$\sqrt{32} \quad \sqrt{12} \quad \sqrt{27} \quad \sqrt{18} \quad \sqrt{23} \quad \sqrt{50}$$



- Between which two integers will $\sqrt{45}$ be located? Explain how you know.

Lesson 3: Existence and Uniqueness of Square and Cube Roots

Classwork

Opening

The numbers in each column are related. Your goal is to determine how they are related, determine which numbers belong in the blank parts of the columns, and write an explanation for how you know the numbers belong there.

Find the Rule Part 1

1	1
2	
3	9
	81
11	121
15	
	49
10	
12	
	169
m	
	n

Find the Rule Part 2

1	1
2	
3	27
	125
6	216
11	
	64
10	
7	
	2,744
p	
	q

Exercises 1–9

Find the positive value of x that makes each equation true. Check your solution.

1. $x^2 = 169$

a. Explain the first step in solving this equation.

b. Solve the equation and check your answer.

2. A square-shaped park has an area of 324 ft^2 . What are the dimensions of the park? Write and solve an equation.

3. $625 = x^2$

4. A cube has a volume of 27 in^3 . What is the measure of one of its sides? Write and solve an equation.

5. What positive value of x makes the following equation true: $x^2 = 64$? Explain.

6. What positive value of x makes the following equation true: $x^3 = 64$? Explain.

7. $x^2 = 256^{-1}$ Find the positive value of x that makes the equation true.

8. $x^3 = 343^{-1}$ Find the positive value of x that makes the equation true.

9. Is 6 a solution to the equation $x^2 - 4 = 5x$? Explain why or why not.

Lesson Summary

The symbol $\sqrt[n]{\quad}$ is called a radical. Then an equation that contains that symbol is referred to as a radical equation. So far we have only worked with square roots ($n = 2$). Technically, we would denote a positive square root as $\sqrt[2]{\quad}$, but it is understood that the symbol $\sqrt{\quad}$ alone represents a positive square root.

When $n = 3$, then the symbol $\sqrt[3]{\quad}$ is used to denote the cube root of a number. Since $x^3 = x \cdot x \cdot x$, then the cube root of x^3 is x , i.e., $\sqrt[3]{x^3} = x$.

The square or cube root of a positive number exists, and there can be only one positive square root or one cube root of the number.

Problem Set

Find the positive value of x that makes each equation true. Check your solution.

1. What positive value of x makes the following equation true: $x^2 = 289$? Explain.
2. A square shaped park has an area of 400 ft^2 . What are the dimensions of the park? Write and solve an equation.
3. A cube has a volume of 64 in^3 . What is the measure of one of its sides? Write and solve an equation.
4. What positive value of x makes the following equation true: $125 = x^3$? Explain.
5. $x^2 = 441^{-1}$ Find the positive value of x that makes the equation true.
 - a. Explain the first step in solving this equation.
 - b. Solve and check your solution.
6. $x^3 = 125^{-1}$ Find the positive value of x that makes the equation true.
7. The area of a square is 196 in^2 . What is the length of one side of the square? Write and solve an equation, then check your solution.
8. The volume of a cube is 729 cm^3 . What is the length of one side of the cube? Write and solve an equation, then check your solution.
9. What positive value of x would make the following equation true: $19 + x^2 = 68$?

Lesson 4: Simplifying Square Roots

Classwork

Opening Exercises 1–6

- What does $\sqrt{16}$ equal?
 - What does 4×4 equal?
 - Does $\sqrt{16} = \sqrt{4 \times 4}$?
- What does $\sqrt{36}$ equal?
 - What does 6×6 equal?
 - Does $\sqrt{36} = \sqrt{6 \times 6}$?
- What does $\sqrt{121}$ equal?
 - What does 11×11 equal?
 - Does $\sqrt{121} = \sqrt{11 \times 11}$?
- What does $\sqrt{81}$ equal?
 - What does 9×9 equal?
 - Does $\sqrt{81} = \sqrt{9 \times 9}$?
- What is another way to write $\sqrt{20}$?
- What is another way to write $\sqrt{28}$?

Example 1

Simplify the square root as much as possible.

$$\sqrt{50} =$$

Example 2

Simplify the square root as much as possible.

$$\sqrt{28} =$$

Exercises 7–10

Simplify the square roots as much as possible.

7. $\sqrt{18}$

8. $\sqrt{44}$

9. $\sqrt{169}$

10. $\sqrt{75}$

Example 3

Simplify the square root as much as possible.

$$\sqrt{128} =$$

Example 4

Simplify the square root as much as possible.

$$\sqrt{288} =$$

Exercises 11–14

11. Simplify $\sqrt{108}$.

12. Simplify $\sqrt{250}$.

13. Simplify $\sqrt{200}$.

14. Simplify $\sqrt{504}$.

Lesson Summary

Square roots of non-perfect squares can be simplified by using the factors of the number. Any perfect square factors of a number can be simplified.


For example:

$$\begin{aligned}\sqrt{72} &= \sqrt{36 \times 2} \\ &= \sqrt{36} \times \sqrt{2} \\ &= \sqrt{6^2} \times \sqrt{2} \\ &= 6 \times \sqrt{2} \\ &= 6\sqrt{2}\end{aligned}$$


Problem Set

Simplify each of the square roots in Problems 1–5 as much as possible.


- $\sqrt{98}$
- $\sqrt{54}$
- $\sqrt{144}$
- $\sqrt{512}$
- $\sqrt{756}$
- What is the length of the unknown side of the right triangle? Simplify your answer.



- What is the length of the unknown side of the right triangle? Simplify your answer.



8. What is the length of the unknown side of the right triangle? Simplify your answer.



9. Josue simplified $\sqrt{450}$ as $15\sqrt{2}$. Is he correct? Explain why or why not.
10. Tiah was absent from school the day that you learned how to simplify a square root. Using $\sqrt{360}$, write Tiah an explanation for simplifying square roots.

Lesson 5: Solving Radical Equations

Classwork

Example 1

Find the positive value of x that makes the equation true.

$$x^3 + 9x = \frac{1}{2}(18x + 54)$$

Example 2

Find the positive value of x that makes the equation true.

$$x(x - 3) - 51 = -3x + 13$$

Exercises 1–8

Find the positive value of x that makes each equation true, and then verify your solution is correct.

1. Solve $x^2 - 14 = 5x + 67 - 5x$.

Explain how you solved the equation.

2. Solve and simplify: $x(x - 1) = 121 - x$.


3. A square has a side length of $3x$ and an area of 324 in^2 . What is the value of x ?

4. $-3x^3 + 14 = -67$

5. $x(x + 4) - 3 = 4(x + 19.5)$

6. $216 + x = x(x^2 - 5) + 6x$

7. What are we trying to determine in the diagram below?



Determine the value of x and check your answer.

Lesson Summary

Equations that contain variables that are squared or cubed can be solved using the properties of equality and the definition of square and cube roots.

Simplify an equation until it is in the form of $x^2 = p$ or $x^3 = p$ where p is a positive rational number, then take the square or cube root to determine the positive value of x .

Example:

Solve for x .

$$\begin{aligned}\frac{1}{2}(2x^2 + 10) &= 30 \\ x^2 + 5 &= 30 \\ x^2 + 5 - 5 &= 30 - 5 \\ x^2 &= 25 \\ \sqrt{x^2} &= \sqrt{25} \\ x &= 5\end{aligned}$$

Check:


$$\begin{aligned}\frac{1}{2}(2(5)^2 + 10) &= 30 \\ \frac{1}{2}(2(25) + 10) &= 30 \\ \frac{1}{2}(50 + 10) &= 30 \\ \frac{1}{2}(60) &= 30 \\ 30 &= 30\end{aligned}$$

Problem Set


Find the positive value of x that makes each equation true, and then verify your solution is correct.

- $x^2(x + 7) = \frac{1}{2}(14x^2 + 16)$
- $x^3 = 1,331^{-1}$
- $\frac{x^9}{x^7} - 49 = 0$. Determine the positive value of x that makes the equation true, and then explain how you solved the equation.
- $(8x)^2 = 1$. Determine the positive value of x that makes the equation true.
- $(9\sqrt{x})^2 - 43x = 76$


6. Determine the length of the hypotenuse of the right triangle below.



7. Determine the length of the legs in the right triangle below.



8. An equilateral triangle has side lengths of 6 cm. What is the height of the triangle? What is the area of the triangle?



9. Challenge: $\left(\frac{1}{2}x\right)^2 - 3x = 7x + 8 - 10x$. Find the positive value of x that makes the equation true.

10. Challenge: $11x + x(x - 4) = 7(x + 9)$. Find the positive value of x that makes the equation true.

Lesson 6: Finite and Infinite Decimals

Classwork

Exercises 1–5

1. Use long division to determine the decimal expansion of $\frac{54}{20}$.

2. Use long division to determine the decimal expansion of $\frac{7}{8}$.

3. Use long division to determine the decimal expansion of $\frac{8}{9}$.

4. Use long division to determine the decimal expansion of $\frac{22}{7}$.

5. What do you notice about the decimal expansions of Exercises 1 and 2 compared to the decimal expansions of Exercises 3 and 4?

Example 1

Consider the fraction $\frac{5}{8}$. Is it equal to a finite decimal? How do you know?

Example 2

Consider the fraction $\frac{17}{125}$. Is it equal to a finite or infinite decimal? How do you know?

Exercises 6–10

Show your steps, but use a calculator for the multiplications.

6. Convert the fraction $\frac{7}{8}$ to a decimal.

a. Write the denominator as a product of 2's or 5's. Explain why this way of rewriting the denominator helps to find the decimal representation of $\frac{7}{8}$.

b. Find the decimal representation of $\frac{7}{8}$. Explain why your answer is reasonable.

7. Convert the fraction $\frac{43}{64}$ to a decimal.

8. Convert the fraction $\frac{29}{125}$ to a decimal.

9. Convert the fraction $\frac{19}{34}$ to a decimal.
10. Identify the type of decimal expansion for each of the numbers in Exercises 6–9 as finite or infinite. Explain why their decimal expansion is such.

Example 3

Write $\frac{7}{80}$ as a decimal. Will it be finite or infinite? Explain.

Example 4

Write $\frac{3}{160}$ as a decimal. Will it be finite or infinite? Explain.

Exercises 11–13

Show your steps, but use a calculator for the multiplications.

11. Convert the fraction $\frac{37}{40}$ to a decimal.

a. Write the denominator as a product of 2's and/or 5's. Explain why this way of rewriting the denominator helps to find the decimal representation of $\frac{37}{40}$.

b. Find the decimal representation of $\frac{37}{40}$. Explain why your answer is reasonable.

12. Convert the fraction $\frac{3}{250}$ to a decimal.

13. Convert the fraction $\frac{7}{1,250}$ to a decimal.

Lesson Summary

Fractions with denominators that can be expressed as products of 2's and/or 5's have decimal expansions that are finite.

Example:

Does the fraction $\frac{1}{8}$ have a finite or infinite decimal expansion?

Since $8 = 2^3$, then the fraction has a finite decimal expansion. The decimal expansion is found by:

$$\frac{1}{8} = \frac{1}{2^3} = \frac{1 \times 5^3}{2^3 \times 5^3} = \frac{125}{10^3} = 0.125$$

When the denominator of a fraction cannot be expressed as a product of 2's and/or 5's then the decimal expansion of the number will be infinite.

When infinite decimals repeat, such as $0.8888888 \dots$ or $0.454545454545 \dots$, they are typically abbreviated using the notation $0.\overline{8}$ and $0.\overline{45}$, respectively. The notation indicates that the digit 8 repeats indefinitely and that the two-digit block 45 repeats indefinitely.

Problem Set

Convert each fraction to a finite decimal. If the fraction cannot be written as a finite decimal, then state how you know. Show your steps, but use a calculator for the multiplications.

1. $\frac{2}{32}$

2. $\frac{99}{125}$

a. Write the denominator as a product of 2's and/or 5's. Explain why this way of rewriting the denominator helps to find the decimal representation of $\frac{99}{125}$.

b. Find the decimal representation of $\frac{99}{125}$. Explain why your answer is reasonable.

3. $\frac{15}{128}$

4. $\frac{8}{15}$

5. $\frac{3}{28}$

6. $\frac{13}{400}$

7. $\frac{5}{64}$

8. $\frac{15}{35}$

9. $\frac{199}{250}$

10. $\frac{219}{625}$

Lesson 7: Infinite Decimals


Classwork

Opening Exercises 1–4

1. Write the expanded form of the decimal 0.3765 using powers of 10.
2. Write the expanded form of the decimal 0.3333333 ... using powers of 10.
3. What is an infinite decimal? Give an example.
4. Do you think it is acceptable to write that $1 = 0.99999 \dots$? Why or why not?


Example 1

The number 0.253 on the number line:



Example 2


The number $\frac{5}{6} = 0.833333 \dots = 0.8\bar{3}$ on the number line:



Exercises 5–10

5. a. Write the expanded form of the decimal 0.125 using powers of 10.


b. Show on the number line the representation of the decimal 0.125.



c. Is the decimal finite or infinite? How do you know?

6. a. Write the expanded form of the decimal 0.3875 using powers of 10.


b. Show on the number line the representation of the decimal 0.3875.



c. Is the decimal finite or infinite? How do you know?

7. a. Write the expanded form of the decimal $0.777777 \dots$ using powers of 10.


b. Show on the number line the representation of the decimal $0.777777 \dots$



c. Is the decimal finite or infinite? How do you know?

8. a. Write the expanded form of the decimal $0.\overline{45}$ using powers of 10.

b. Show on the number line the representation of the decimal $0.\overline{45}$.



- c. Is the decimal finite or infinite? How do you know?
9. Order the following numbers from least to greatest: 2.121212 , 2.1 , 2.2 , and $2.\overline{12}$.
10. Explain how you knew which order to put the numbers in.


Lesson Summary

An infinite decimal is a decimal whose expanded form and number line representation are infinite.

Example:

The expanded form of the decimal $0.8\overline{3}$ is $0.8\overline{3} = \frac{8}{10} + \frac{3}{10^2} + \frac{3}{10^3} + \frac{3}{10^4} + \dots$

The number is represented on the number line shown below. Each new line is a magnification of the interval shown above it. For example, the first line is the unit from 0 to 1 divided into 10 equal parts, or tenths. The second line is the interval from 0.8 to 0.9 divided into ten equal parts, or hundredths. The third line is the interval from 0.83 to 0.84 divided into ten equal parts, or thousandths, and so on.




With each new line we are representing an increasingly smaller value of the number, so small that the amount approaches a value of 0. Consider the 20th line of the picture above. We would be adding $\frac{3}{10^{20}}$ to the value of the number, which is 0.00000000000000000003. It should be clear that $\frac{3}{10^{20}}$ is a very small number and is fairly close to a value of 0.


This reasoning is what we use to explain why the value of the infinite decimal $0.\overline{9}$ is 1.

Problem Set

1. a. Write the expanded form of the decimal 0.625 using powers of 10.
- b. Show the representation of the decimal 0.625 on the number line.
- c. Is the decimal finite or infinite? How do you know?



2. a. Write the expanded form of the decimal $0.\overline{370}$ using powers of 10.
- b. Show on the number line the representation of the decimal 0.370370 ...
- c. Is the decimal finite or infinite? How do you know?



3. Which is a more accurate representation of the number $\frac{2}{3}$: 0.6666 or $0.\overline{6}$? Explain. Which would you prefer to compute with?
4. Explain why we shorten infinite decimals to finite decimals to perform operations. Explain the effect of shortening an infinite decimal on our answers.
5. A classmate missed the discussion about why $0.\overline{9} = 1$. Convince your classmate that this equality is true.
6. Explain why $0.3333 < 0.\overline{3}$.

Lesson 8: The Long Division Algorithm

Classwork

Example 1

Show that the decimal expansion of $\frac{26}{4}$ is 6.5.

Exercises 1–5

1. Use long division to determine the decimal expansion of $\frac{142}{2}$.

a. Fill in the blanks to show another way to determine the decimal expansion of $\frac{142}{2}$.

$$142 = \underline{\quad} \times 2 + \underline{\quad}$$

$$\frac{142}{2} = \frac{\underline{\quad} \times 2 + \underline{\quad}}{2}$$

$$\frac{142}{2} = \frac{\underline{\quad} \times 2}{2} + \frac{\underline{\quad}}{2}$$

$$\frac{142}{2} = \underline{\quad} + \frac{\underline{\quad}}{2}$$

$$\frac{142}{2} = \underline{\quad}$$

b. Does the number $\frac{142}{2}$ have a finite or infinite decimal expansion? Explain how you know.

2. Use long division to determine the decimal expansion of $\frac{142}{4}$.

a. Fill in the blanks to show another way to determine the decimal expansion of $\frac{142}{4}$.

$$142 = \underline{\quad} \times 4 + \underline{\quad}$$

$$\frac{142}{4} = \frac{\underline{\quad} \times 4 + \underline{\quad}}{4}$$

$$\frac{142}{4} = \frac{\underline{\quad}}{4} + \frac{\underline{\quad}}{4}$$

$$\frac{142}{4} = \underline{\quad} + \frac{\underline{\quad}}{4}$$

$$\frac{142}{4} = \underline{\quad}$$

b. Does the number $\frac{142}{4}$ have a finite or infinite decimal expansion? Explain how you know.

3. Use long division to determine the decimal expansion of $\frac{142}{6}$.

- a. Fill in the blanks to show another way to determine the decimal expansion of $\frac{142}{6}$.

$$142 = \underline{\quad} \times 6 + \underline{\quad}$$

$$\frac{142}{6} = \frac{\underline{\quad} \times 6 + \underline{\quad}}{6}$$

$$\frac{142}{6} = \frac{\underline{\quad} \times 6}{6} + \frac{\underline{\quad}}{6}$$

$$\frac{142}{6} = \underline{\quad} + \frac{\underline{\quad}}{6}$$

$$\frac{142}{6} = \underline{\hspace{2cm}}$$

- b. Does the number $\frac{142}{6}$ have a finite or infinite decimal expansion? Explain how you know.

4. Use long division to determine the decimal expansion of $\frac{142}{11}$.

- a. Fill in the blanks to show another way to determine the decimal expansion of $\frac{142}{11}$.

$$142 = \underline{\quad} \times 11 + \underline{\quad}$$

$$\frac{142}{11} = \frac{\underline{\quad} \times 11 + \underline{\quad}}{11}$$

$$\frac{142}{11} = \frac{\underline{\quad} \times 11}{11} + \frac{\underline{\quad}}{11}$$

$$\frac{142}{11} = \underline{\quad} + \frac{\underline{\quad}}{11}$$

$$\frac{142}{11} = \underline{\hspace{2cm}}$$

- b. Does the number $\frac{142}{11}$ have a finite or infinite decimal expansion? Explain how you know.

5. Which fractions produced an infinite decimal expansion? Why do you think that is?

Exercises 6–10

6. Does the number $\frac{65}{13}$ have a finite or infinite decimal expansion? Based on our definition of rational numbers having a decimal expansion that repeats eventually, is the number rational? Explain.

7. Does the number $\frac{17}{11}$ have a finite or infinite decimal expansion? Based on our definition of rational numbers having a decimal expansion that repeats eventually, is the number rational? Explain.
8. Does the number $\pi = 3.1415926535897 \dots$ have a finite or infinite decimal expansion? Based on our definition of rational numbers having a decimal expansion that repeats eventually, is the number rational? Explain.
9. Does the number $\frac{860}{999} = 0.860860860 \dots$ have a finite or infinite decimal expansion? Based on our definition of rational numbers having a decimal expansion that repeats eventually, is the number rational? Explain.
10. Does the number $\sqrt{2} = 1.41421356237 \dots$ have a finite or infinite decimal expansion? Based on our definition of rational numbers having a decimal expansion that repeats eventually, is the number rational? Explain.

Lesson Summary

The long division algorithm is a procedure that can be used to determine the decimal expansion of infinite decimals. Every rational number has a decimal expansion that repeats eventually. For example, the number 32 is rational because it has a repeat block of the digit 0 in its decimal expansion, $32.\bar{0}$. The number $\frac{1}{3}$ is rational because it has a repeat block of the digit 3 in its decimal expansion, $0.\bar{3}$. The number $0.454545 \dots$ is rational because it has a repeat block of the digits 45 in its decimal expansion, $0.\overline{45}$.

Problem Set

1. Write the decimal expansion of $\frac{7000}{9}$. Based on our definition of rational numbers having a decimal expansion that repeats eventually, is the number rational? Explain.
2. Write the decimal expansion of $\frac{6555555}{3}$. Based on our definition of rational numbers having a decimal expansion that repeats eventually, is the number rational? Explain.
3. Write the decimal expansion of $\frac{350000}{11}$. Based on our definition of rational numbers having a decimal expansion that repeats eventually, is the number rational? Explain.
4. Write the decimal expansion of $\frac{12000000}{37}$. Based on our definition of rational numbers having a decimal expansion that repeats eventually, is the number rational? Explain.
5. Someone notices that the long division of 2,222,222 by 6 has a quotient of 370,370 and remainder 2 and wonders why there is a repeating block of digits in the quotient, namely 370. Explain to the person why this happens.
6. Is the number $\frac{9}{11} = 0.81818181 \dots$ rational? Explain.
7. Is the number $\sqrt{3} = 1.73205080 \dots$ rational? Explain.
8. Is the number $\frac{41}{333} = 0.1231231231 \dots$ rational? Explain.

Example 2

Write the decimal expansion of the fraction $\frac{17}{125}$.

Example 3

Write the decimal expansion of the fraction $\frac{35}{11}$.

Example 4

Write the decimal expansion of the fraction $\frac{6}{7}$.

Exercises 3–5

3. a. Choose a power of ten to use to convert this fraction to a decimal: $\frac{4}{13}$. Explain your choice.

b. Determine the decimal expansion of $\frac{4}{13}$ and verify you are correct using a calculator.

4. Write the decimal expansion of $\frac{1}{11}$. Verify you are correct using a calculator.

5. Write the decimal expansion of $\frac{19}{21}$. Verify you are correct using a calculator.

Lesson Summary

Multiplying a fraction's numerator and denominator by the same power of 10 to determine its decimal expansion is similar to including extra zeroes to the right of a decimal when using the long division algorithm. The method of multiplying by a power of 10 reduces the work to whole number division.

Example: We know that the fraction $\frac{5}{3}$ has an infinite decimal expansion because the denominator is not a product of 2's and/or 5's. Its decimal expansion is found by the following procedure:

$$\begin{aligned} \frac{5}{3} &= \frac{5 \times 10^2}{3} \times \frac{1}{10^2} && \text{Multiply numerator and denominator by } 10^2 \\ &= \frac{166 \times 3 + 2}{3} \times \frac{1}{10^2} && \text{Rewrite the numerator as a product of a number multiplied by the denominator} \\ &= \left(\frac{166 \times 3}{3} + \frac{2}{3} \right) \times \frac{1}{10^2} && \text{Rewrite the first term as a sum of fractions with the same denominator} \\ &= \left(166 + \frac{2}{3} \right) \times \frac{1}{10^2} && \text{Simplify} \\ &= \frac{166}{10^2} + \left(\frac{2}{3} \times \frac{1}{10^2} \right) && \text{Use the distributive property} \\ &= 1.66 + \left(\frac{2}{3} \times \frac{1}{10^2} \right) && \text{Simplify} \\ &= 166 \times \frac{1}{10^2} + \frac{2}{3} \times \frac{1}{10^2} && \text{Simplify the first term using what you know about place value} \end{aligned}$$

Notice that the value of the remainder, $\left(\frac{2}{3} \times \frac{1}{10^2} \right) = \frac{2}{300} = 0.006$, is quite small and does not add much value to the number. Therefore, 1.66 is a good estimate of the value of the infinite decimal for the fraction $\frac{5}{3}$.

Problem Set

- Choose a power of ten to convert this fraction to a decimal: $\frac{4}{11}$. Explain your choice.
 - Determine the decimal expansion of $\frac{4}{11}$ and verify you are correct using a calculator.
- Write the decimal expansion of $\frac{5}{13}$. Verify you are correct using a calculator.
- Write the decimal expansion of $\frac{23}{39}$. Verify you are correct using a calculator.

4. Tamer wrote the decimal expansion of $\frac{3}{7}$ as 0.418571, but when he checked it on a calculator it was 0.428571. Identify his error and explain what he did wrong.

$$\begin{aligned}\frac{3}{7} &= \frac{3 \times 10^6}{7} \times \frac{1}{10^6} \\ &= \frac{3,000,000}{7} \times \frac{1}{10^6}\end{aligned}$$

$$3,000,000 = 418,571 \times 7 + 3$$

$$\begin{aligned}\frac{3}{7} &= \frac{418571 \times 7 + 3}{7} \times \frac{1}{10^6} \\ &= \left(\frac{418571 \times 7}{7} + \frac{3}{7} \right) \times \frac{1}{10^6} \\ &= \left(418571 + \frac{3}{7} \right) \times \frac{1}{10^6} \\ &= 418,571 \times \frac{1}{10^6} + \left(\frac{3}{7} \times \frac{1}{10^6} \right) \\ &= \frac{418571}{10^6} + \left(\frac{3}{7} \times \frac{1}{10^6} \right) \\ &= 0.418571 + \left(\frac{3}{7} \times \frac{1}{10^6} \right)\end{aligned}$$

5. Given that $\frac{6}{7} = 0.857142 + \left(\frac{6}{7} \times \frac{1}{10^6} \right)$. Explain why 0.857142 is a good estimate of $\frac{6}{7}$.

Lesson 10: Converting Repeating Decimals to Fractions

Classwork

Example 1

Find the fraction that is equal to the infinite decimal $0.\overline{81}$.

Exercises 1–2

- Let $x = 0.\overline{123}$. Explain why multiplying both sides of this equation by 10^3 will help us determine the fractional representation of x .

- b. After multiplying both sides of the equation by 10^3 , rewrite the resulting equation by making a substitution that will help determine the fractional value of x . Explain how you were able to make the substitution.
- c. Solve the equation to determine the value of x .
- d. Is your answer reasonable? Check your answer using a calculator.

2. Find the fraction equal to $0.\overline{4}$. Check that you are correct using a calculator.

Example 2

Find the fraction that is equal to the infinite decimal $2.13\overline{8}$.

Exercises 3–4

3. Find the fraction equal to $1.\overline{623}$. Check that you are correct using a calculator.

4. Find the fraction equal to $2.\overline{960}$. Check that you are correct using a calculator.

Lesson Summary

Numbers with decimal expansions that repeat are rational numbers and can be converted to fractions using a linear equation.

Example: Find the fraction that is equal to the number $0.\overline{567}$.

Let x represent the infinite decimal $0.\overline{567}$.

$x = 0.\overline{567}$	
$10^3x = 10^3(0.\overline{567})$	Multiply by 10^3 because there are 3 digits that repeat
$1000x = 567.\overline{567}$	Simplify
$1000x = 567 + 0.\overline{567}$	By addition
$1000x = 567 + x$	By substitution; $x = 0.\overline{567}$
$1000x - x = 567 + x - x$	Subtraction Property of Equality
$999x = 567$	Simplify
$\frac{999}{999}x = \frac{567}{999}$	Division Property of Equality
$x = \frac{567}{999} = \frac{63}{111}$	Simplify

This process may need to be used more than once when the repeating digits do not begin immediately after the decimal. For numbers such as $1.2\overline{6}$, for example.

Irrational numbers are numbers that are not rational. They have infinite decimals that do not repeat and cannot be represented as a fraction.

Problem Set

1.
 - a. Let $x = 0.\overline{631}$. Explain why multiplying both sides of this equation by 10^3 will help us determine the fractional representation of x .
 - b. After multiplying both sides of the equation by 10^3 , rewrite the resulting equation by making a substitution that will help determine the fractional value of x . Explain how you were able to make the substitution.
 - c. Solve the equation to determine the value of x .
 - d. Is your answer reasonable? Check your answer using a calculator.
2. Find the fraction equal to $3.40\overline{8}$. Check that you are correct using a calculator.
3. Find the fraction equal to $0.\overline{5923}$. Check that you are correct using a calculator.
4. Find the fraction equal to $2.3\overline{82}$. Check that you are correct using a calculator.
5. Find the fraction equal to $0.\overline{714285}$. Check that you are correct using a calculator.

6. Explain why an infinite decimal that is not a repeating decimal cannot be rational.
7. In a previous lesson we were convinced that it is acceptable to write $0.\overline{9} = 1$. Use what you learned today to show that it is true.
8. Examine the following repeating infinite decimals and their fraction equivalents. What do you notice? Why do you think what you observed is true?

$$0.\overline{81} = \frac{81}{99}$$

$$0.\overline{4} = \frac{4}{9}$$

$$0.\overline{123} = \frac{123}{999}$$

$$0.\overline{60} = \frac{60}{99}$$

$$0.\overline{9} = 1.0$$

Lesson 11: The Decimal Expansion of Some Irrational Numbers

Classwork

Opening Exercise

Place $\sqrt{28}$ on a number line. What decimal do you think $\sqrt{28}$ is equal to? Explain your reasoning.

Example 1


Recall the Basic Inequality:

Let c and d be two positive numbers, and let n be a fixed positive integer. Then $c < d$ if and only if $c^n < d^n$.


Write the decimal expansion of $\sqrt{3}$.

First approximation:

Second approximation:



Third approximation:




Example 2


Write the decimal expansion of $\sqrt{28}$.

First approximation:


Second approximation:



Third approximation:



Fourth approximation:



Exercise 2

Between which interval of hundredths would $\sqrt{14}$ be located? Show your work.

Lesson Summary

To get the decimal expansion of a square root of a non-perfect square you must use the method of rational approximation. Rational approximation is a method that uses a sequence of rational numbers to get closer and closer to a given number to estimate the value of the number. The method requires that you investigate the size of the number by examining its value for increasingly smaller powers of 10 (i.e., tenths, hundredths, thousandths, and so on). Since $\sqrt{22}$ is not a perfect square, you would use rational approximation to determine its decimal expansion.

Example:

Begin by determining which two integers the number would lie.

$\sqrt{22}$ is between the integers 4 and 5 because $4^2 < (\sqrt{22})^2 < 5^2$, which is equal to $16 < 22 < 25$.

Next, determine which interval of tenths the number belongs.

$\sqrt{22}$ is between 4.6 and 4.7 because $4.6^2 < (\sqrt{22})^2 < 4.7^2$, which is equal to $21.16 < 22 < 22.09$.

Next, determine which interval of hundredths the number belongs.

$\sqrt{22}$ is between 4.69 and 4.70 because $4.69^2 < (\sqrt{22})^2 < 4.70^2$, which is equal to $21.9961 < 22 < 22.09$.

A good estimate of the value of $\sqrt{22}$ is 4.69 because 22 is closer to 21.9961 than it is to 22.09.

Notice that with each step we are getting closer and closer to the actual value, 22. This process can continue using intervals of thousandths, ten-thousandths, and so on.

Any number that cannot be expressed as a rational number is called an *irrational number*. Irrational numbers are those numbers with decimal expansions that are infinite and do not have a repeating block of digits.

Problem Set

- Use the method of rational approximation to determine the decimal expansion of $\sqrt{84}$. Determine which interval of hundredths it would lie in.
- Get a 3 decimal digit approximation of the number $\sqrt{34}$.
- Write the decimal expansion of $\sqrt{47}$ to at least 2 decimal digits.
- Write the decimal expansion of $\sqrt{46}$ to at least 2 decimal digits.
- Explain how to improve the accuracy of decimal expansion of an irrational number.
- Is the number $\sqrt{125}$ rational or irrational? Explain.
- Is the number 0.646464646 ... rational or irrational? Explain.
- Is the number 3.741657387 ... rational or irrational? Explain.
- Is the number $\sqrt{99}$ rational or irrational? Explain.
- Challenge: Get a 2 decimal digit approximation of the number $\sqrt[3]{9}$.

Lesson 12: Decimal Expansions of Fractions, Part 2

Classwork

Example 1

Write the decimal expansion of $\frac{35}{11}$.

Exercises 1–3

1. Use rational approximation to determine the decimal expansion of $\frac{5}{3}$.

2. Use rational approximation to determine the decimal expansion of $\frac{5}{11}$.

3. a. Determine the decimal expansion of the number $\frac{23}{99}$ using rational approximation and long division.

b. When comparing rational approximation to long division, what do you notice?

Lesson Summary

The method of rational approximation, used earlier to write the decimal expansion of irrational numbers, can also be used to write the decimal expansion of fractions (rational numbers).

When used with rational numbers, there is no need to guess and check to determine the interval of tenths, hundredths, thousandths, etc. in which a number will lie. Rather, computation can be used to determine between which two consecutive integers, m and $m + 1$, a number would lie for a given place value. For example, to determine where the fraction $\frac{1}{8}$ lies in the interval of tenths, compute using the following inequality:

$$\begin{array}{ll} \frac{m}{10} < \frac{1}{8} < \frac{m+1}{10} & \text{Use the denominator of 10 because of our need to find the tenths digit of } \frac{1}{8} \\ m < \frac{10}{8} < m+1 & \text{Multiply through by 10} \\ m < 1\frac{1}{4} < m+1 & \text{Simplify the fraction } \frac{10}{8} \end{array}$$

The last inequality implies that $m = 1$ and $m + 1 = 2$, because $1 < 1\frac{1}{4} < 2$. Then the tenths digit of the decimal expansion of $\frac{1}{8}$ is 1.

Next, find the difference between the number $\frac{1}{8}$ and the known tenths digit value, $\frac{1}{10}$, i.e., $\frac{1}{8} - \frac{1}{10} = \frac{2}{80} = \frac{1}{40}$.

Use the inequality again, this time with $\frac{1}{40}$, to determine the hundredths digit of the decimal expansion of $\frac{1}{8}$.

$$\begin{array}{ll} \frac{m}{100} < \frac{1}{40} < \frac{m+1}{100} & \text{Use the denominator of 100 because of our need to find the hundredths digit of } \frac{1}{8} \\ m < \frac{100}{40} < m+1 & \text{Multiply through by 100} \\ m < 2\frac{1}{2} < m+1 & \text{Simplify the fraction } \frac{100}{40} \end{array}$$

The last inequality implies that $m = 2$ and $m + 1 = 3$, because $2 < 2\frac{1}{2} < 3$. Then the hundredths digit of the decimal expansion of $\frac{1}{8}$ is 2.

Continue the process until the decimal expansion is complete or you notice a pattern of repeating digits.

Problem Set

1. Explain why the tenths digit of $\frac{3}{11}$ is 2, using rational approximation.
2. Use rational approximation to determine the decimal expansion of $\frac{25}{9}$.
3. Use rational approximation to determine the decimal expansion of $\frac{11}{41}$ to at least 5 digits.
4. Use rational approximation to determine which number is larger, $\sqrt{10}$ or $\frac{28}{9}$.
5. Sam says that $\frac{7}{11} = 0.63$, and Jaylen says that $\frac{7}{11} = 0.636$. Who is correct? Why?

Lesson 13: Comparison of Irrational Numbers

Classwork

Exercises 1–11

1. Rodney thinks that $\sqrt[3]{64}$ is greater than $\frac{17}{4}$. Sam thinks that $\frac{17}{4}$ is greater. Who is right and why?

2. Which number is smaller, $\sqrt[3]{27}$ or 2.89? Explain.

3. Which number is smaller, $\sqrt{121}$ or $\sqrt[3]{125}$? Explain.

4. Which number is smaller, $\sqrt{49}$ or $\sqrt[3]{216}$? Explain.


5. Which number is greater, $\sqrt{50}$ or $\frac{319}{45}$? Explain.

6. Which number is greater, $\frac{5}{11}$ or $0.\bar{4}$? Explain.

7. Which number is greater, $\sqrt{38}$ or $\frac{154}{25}$? Explain.


8. Which number is greater, $\sqrt{2}$ or $\frac{15}{9}$? Explain.

9. Place the following numbers at their approximate location on the number line: $\sqrt{25}$, $\sqrt{28}$, $\sqrt{30}$, $\sqrt{32}$, $\sqrt{35}$, $\sqrt{36}$.



10. Challenge: Which number is larger $\sqrt{5}$ or $\sqrt[3]{11}$?

11. A certain chessboard is being designed so that each square has an area of 3 in^2 . What is the length, rounded to the tenths place, of one edge of the board? (A chessboard is composed of 64 squares as shown.)




Lesson Summary

The decimal expansion of rational numbers can be found by using long division, equivalent fractions, or the method of rational approximation.


The decimal expansion of irrational numbers can be found using the method of rational approximation.

Problem Set

- Which number is smaller, $\sqrt[3]{343}$ or $\sqrt{48}$? Explain.
- Which number is smaller, $\sqrt{100}$ or $\sqrt[3]{1000}$? Explain.
- Which number is larger, $\sqrt{87}$ or $\frac{929}{99}$? Explain.
- Which number is larger, $\frac{9}{13}$ or $0.\overline{692}$? Explain.
- Which number is larger, 9.1 or $\sqrt{82}$? Explain.
- Place the following numbers at their approximate location on the number line: $\sqrt{144}$, $\sqrt[3]{1000}$, $\sqrt{130}$, $\sqrt{110}$, $\sqrt{120}$, $\sqrt{115}$, $\sqrt{133}$. Explain how you knew where to place the numbers.



- Which of the two right triangles shown below, measured in units, has the longer hypotenuse? Approximately how much longer is it?




Lesson 14: The Decimal Expansion of π


Classwork

Opening Exercises 1–3


1. Write an equation for the area, A , of the circle shown.



2. Write an equation for the circumference, C , of the circle shown.



3. Each of the squares in the grid below has an area of 1 unit².



- a. Estimate the area of the circle shown by counting squares.
- b. Calculate the area of the circle using a radius of 5 units and 3.14 for π .

Exercises 4–7

4. Gerald and Sarah are building a wheel with a radius of 6.5 cm and are trying to determine the circumference. Gerald says, “Because $6.5 \times 2 \times 3.14 = 40.82$, the circumference is 40.82 cm.” Sarah says, “Because $6.5 \times 2 \times 3.10 = 40.3$ and $6.5 \times 2 \times 3.21 = 41.73$, the circumference is somewhere between 40.3 and 41.73.” Explain the thinking of each student.
5. Estimate the value of the irrational number $(6.12486 \dots)^2$.
6. Estimate the value of the irrational number $(9.204107 \dots)^2$.
7. Estimate the value of the irrational number $(4.014325 \dots)^2$.

Lesson Summary

Irrational numbers, such as π , are frequently approximated in order to compute with them. Common approximations for π are 3.14 and $\frac{22}{7}$. It should be understood that using an approximate value of an irrational number for computations produces an answer that is accurate to only the first few decimal digits.






Problem Set



1. Caitlin estimated π to be $3.10 < \pi < 3.21$. If she uses this approximation of π to determine the area of a circle with a radius of 5 cm, what could the area be?
2. Myka estimated the circumference of a circle with a radius of 4.5 in. to be 28.44 in. What approximate value of π did she use? Is it an acceptable approximation of π ? Explain.
3. A length of ribbon is being cut to decorate a cylindrical cookie jar. The ribbon must be cut to a length that stretches the length of the circumference of the jar. There is only enough ribbon to make one cut. When approximating π to calculate the circumference of the jar, which number in the interval $3.10 < \pi < 3.21$ should be used? Explain.
4. Estimate the value of the irrational number $(1.86211 \dots)^2$.
5. Estimate the value of the irrational number $(5.9035687 \dots)^2$.
6. Estimate the value of the irrational number $(12.30791 \dots)^2$.
7. Estimate the value of the irrational number $(0.6289731 \dots)^2$.
8. Estimate the value of the irrational number $(1.112223333 \dots)^2$.
9. Which number is a better estimate for π , $\frac{22}{7}$ or 3.14? Explain.
10. To how many decimal digits can you correctly estimate the value of the irrational number $(4.56789012 \dots)^2$?

Lesson 15: Pythagorean Theorem, Revisited

Classwork

Proof of the Pythagorean Theorem






Lesson Summary


The Pythagorean Theorem can be proven by showing that the sum of the areas of the squares constructed off of the legs of a right triangle is equal to the area of the square constructed off of the hypotenuse of the right triangle.

Problem Set


1. For the right triangle shown below, identify and use similar triangles to illustrate the Pythagorean Theorem.




2. For the right triangle shown below, identify and use squares formed by the sides of the triangle to illustrate the Pythagorean Theorem.



3. Reese claimed that any figure can be drawn off the sides of a right triangle and that as long as they are similar figures, then the sum of the areas off of the legs will equal the area off of the hypotenuse. She drew the diagram at right by constructing rectangles off of each side of a known right triangle. Is Reese’s claim correct for this example? In order to prove or disprove Reese’s claim, you must first show that the rectangles are similar. If they are, then you can use computations to show that the sum of the areas of the figures off of the sides a and b equal the area of the figure off of side c .



4. After learning the proof of the Pythagorean Theorem using areas of squares, Joseph got really excited and tried explaining it to his younger brother using the diagram to the right. He realized during his explanation that he had done something wrong. Help Joseph find his error. Explain what he did wrong.





5. Draw a right triangle with squares constructed off of each side that Joseph can use the next time he wants to show his younger brother the proof of the Pythagorean Theorem.
6. Explain the meaning of the Pythagorean Theorem in your own words.
7. Draw a diagram that shown an example illustrating the Pythagorean Theorem.

Lesson 16: Converse of the Pythagorean Theorem

Classwork


Proof of the Converse of the Pythagorean Theorem




Exercises 1–7

1. Is the triangle with leg lengths of 3 mi., 8 mi., and hypotenuse of length $\sqrt{73}$ mi. a right triangle? Show your work, and answer in a complete sentence.

2. What is the length of the unknown side of the right triangle shown below? Show your work, and answer in a complete sentence. Provide an exact answer and an approximate answer rounded to the tenths place.




3. What is the length of the unknown side of the right triangle shown below? Show your work, and answer in a complete sentence. Provide an exact answer and an approximate answer rounded to the tenths place.




4. Is the triangle with leg lengths of 9 in., 9 in., and hypotenuse of length $\sqrt{175}$ in. a right triangle? Show your work, and answer in a complete sentence.

5. Is the triangle with leg lengths of $\sqrt{28}$ cm, 6 cm, and hypotenuse of length 8 cm a right triangle? Show your work, and answer in a complete sentence.

6. What is the length of the unknown side of the right triangle shown below? Show your work, and answer in a complete sentence.



7. The triangle shown below is an isosceles right triangle. Determine the length of the legs of the triangle. Show your work, and answer in a complete sentence.




Lesson Summary

The converse of the Pythagorean Theorem states that if a triangle with side lengths a , b , and c satisfies $a^2 + b^2 = c^2$, then the triangle is a right triangle.


The converse can be proven using concepts related to congruence.

Problem Set


1. What is the length of the unknown side of the right triangle shown below? Show your work, and answer in a complete sentence. Provide an exact answer and an approximate answer rounded to the tenths place.



2. What is the length of the unknown side of the right triangle shown below? Show your work, and answer in a complete sentence. Provide an exact answer and an approximate answer rounded to the tenths place.



3. Is the triangle with leg lengths of $\sqrt{3}$ cm, 9 cm, and hypotenuse of length $\sqrt{84}$ cm a right triangle? Show your work, and answer in a complete sentence.
4. Is the triangle with leg lengths of $\sqrt{7}$ km, 5 km, and hypotenuse of length $\sqrt{48}$ km a right triangle? Show your work, and answer in a complete sentence.
5. What is the length of the unknown side of the right triangle shown below? Show your work, and answer in a complete sentence. Provide an exact answer and an approximate answer rounded to the tenths place.



6. Is the triangle with leg lengths of 3, 6, and hypotenuse of length $\sqrt{45}$ a right triangle? Show your work, and answer in a complete sentence.
7. What is the length of the unknown side of the right triangle shown below? Show your work, and answer in a complete sentence. Provide an exact answer and an approximate answer rounded to the tenths place.




8. Is the triangle with leg lengths of 1, $\sqrt{3}$, and hypotenuse of length 2 a right triangle? Show your work, and answer in a complete sentence.
9. Corey found the hypotenuse of a right triangle with leg lengths of 2 and 3 to be $\sqrt{13}$. Corey claims that since $\sqrt{13} = 3.61$ when estimating to two decimal digits, that a triangle with leg lengths of 2, 3, and a hypotenuse of 3.61 is a right triangle. Is he correct? Explain.
10. Explain a proof of the Pythagorean Theorem.
11. Explain a proof of the converse of the Pythagorean Theorem.

Lesson 17: Distance on the Coordinate Plane


Classwork

Example 1


What is the distance between the two points A , B on the coordinate plane?




What is the distance between the two points A , B on the coordinate plane?



What is the distance between the two points A , B on the coordinate plane? Round your answer to the tenths place.




Example 2




Exercises 1–4

For each of the Exercises 1–4, determine the distance between points A and B on the coordinate plane. Round your answer to the tenths place.


1.




2.



3.




4.



Example 3

Is the triangle formed by the points A , B , C a right triangle?



Lesson Summary


To determine the distance between two points on the coordinate plane, begin by connecting the two points. Then draw a vertical line through one of the points and a horizontal line through the other point. The intersection of the vertical and horizontal lines forms a right triangle to which the Pythagorean Theorem can be applied.

To verify if a triangle is a right triangle, use the converse of the Pythagorean Theorem.


Problem Set

For each of the Problems 1–4 determine the distance between points *A* and *B* on the coordinate plane. Round your answer to the tenths place.


1.




2.




3.



4.



5. Is the triangle formed by points A , B , C a right triangle?




Lesson 18: Applications of the Pythagorean Theorem


Classwork

Exercises 1–5

1. The area of the right triangle shown below is 26.46 in^2 . What is the perimeter of the right triangle? Round your answer to the tenths place.




2. The diagram below is a representation of a soccer goal.



- a. Determine the length of the bar, c , that would be needed to provide structure to the goal. Round your answer to the tenths place.
- b. How much netting (in square feet) is needed to cover the entire goal?

3. The typical ratio of length to width that is used to produce televisions is 4:3.



- a. A TV with those exact measurements would be quite small, so generally the size of the television is enlarged by multiplying each number in the ratio by some factor of x . For example, a reasonably sized television might have dimensions of $4 \times 5: 3 \times 5$, where the original ratio 4:3 was enlarged by a scale factor of 5. The size of a television is described in inches, such as a 60" TV, for example. That measurement actually refers to the diagonal length of the TV (distance from an upper corner to the opposite lower corner). What measurement would be applied to a television that was produced using the ratio of $4 \times 5: 3 \times 5$?
- b. A 42" TV was just given to your family. What are the length and width measurements of the TV?
- c. Check that the dimensions you got in part (b) are correct using the Pythagorean Theorem.
- d. The table that your TV currently rests on is 30" in length. Will the new TV fit on the table? Explain.


4. Determine the distance between the following pairs of points. Round your answer to the tenths place. Use graph paper if necessary.

a. $(7, 4)$ and $(-3, -2)$

b. $(-5, 2)$ and $(3, 6)$


c. Challenge: (x_1, y_1) and (x_2, y_2) . Explain your answer.

5. What length of ladder will be needed to reach a height of 7 feet along the wall when the base of the ladder is 4 feet from the wall? Round your answer to the tenths place.




Problem Set

1. A 70" TV is advertised on sale at a local store. What are the length and width of the television?
2. There are two paths that one can use to go from Sarah's house to James' house. One way is to take C Street, and the other way requires you to use A Street and B Street. How much shorter is the direct path along C Street?




3. An isosceles right triangle refers to a right triangle with equal leg lengths, s , as shown below.




What is the length of the hypotenuse of an isosceles right triangle with a leg length of 9 cm? Write an exact answer using a square root and an approximate answer rounded to the tenths place.

4. The area of the right triangle shown at right is 66.5 cm^2 .
- What is the height of the triangle?
 - What is the perimeter of the right triangle? Round your answer to the tenths place.



5. What is the distance between points $(1, 9)$ and $(-4, -1)$? Round your answer to the tenths place.
6. An equilateral triangle is shown below. Determine the area of the triangle. Round your answer to the tenths place.




Lesson 19: Cones and Spheres

Classwork


Opening Exercises 1–2

Note: Figures not drawn to scale.


- Determine the volume for each figure below.
 - Write an expression that shows volume in terms of the area of the base, B , and the height of the figure. Explain the meaning of the expression, and then use it to determine the volume of the figure.




- Write an expression that shows volume in terms of the area of the base, B , and the height of the figure. Explain the meaning of the expression, and then use it to determine the volume of the figure.



2. a. Write an expression that shows volume in terms of the area of the base, B , and the height of the figure. Explain the meaning of the expression, and then use it to determine the volume of the figure.




- b. The volume of the pyramid shown below is 480 in^3 . What do you think the formula to find the volume of a pyramid is? Explain your reasoning.




Example 1


State as many facts as you can about a cone.

**Exercises 3–10**


3. What is the lateral length of the cone shown below?




4. Determine the exact volume of the cone shown below.




5. What is the lateral length (slant height) of the pyramid shown below? Give an exact square root answer and an approximate answer rounded to the tenths place.




6. Determine the volume of the pyramid shown below. Give an exact answer using a square root.




7. What is the length of the chord of the sphere shown below? Give an exact answer using a square root.




8. What is the length of the chord of the sphere shown below? Give an exact answer using a square root.



9. What is the volume of the sphere shown below? Give an exact answer using a square root.




10. What is the volume of the sphere shown below? Give an exact answer using a square root.



Lesson Summary


The volume formula for a right square pyramid is $V = \frac{1}{3}Bh$, where B is the area of the square base.

The lateral length of a cone, sometimes referred to as the slant height, is the side s , shown in the diagram below.



Given the lateral length and the length of the radius, the Pythagorean Theorem can be used to determine the height of the cone.


Let O be the center of a circle, and let P and Q be two points on the circle. Then PQ is called a chord of the circle.




The segments OP and OQ are equal in length because both represent the radius of the circle. If the angle formed by POQ is a right angle, then the Pythagorean Theorem can be used to determine the length of the radius when given the length of the chord; or the length of the chord can be determined if given the length of the radius.

Problem Set


1. What is the lateral length of the cone shown below? Give an approximate answer rounded to the tenths place.




2. What is the volume of the cone shown below? Give an exact answer.



3. Determine the volume and surface area of the pyramid shown below. Give exact answers.




4. Alejandra computed the volume of the cone shown below as $64\pi \text{ cm}^2$. Her work is shown below. Is she correct? If not, explain what she did wrong and calculate the correct volume of the cone. Give an exact answer.




$$\begin{aligned} V &= \frac{1}{3}\pi(4^2)(12) \\ &= \frac{16(12)\pi}{3} \\ &= 64\pi \\ &= 64\pi \text{ cm}^3 \end{aligned}$$

5. What is the length of the chord of the sphere shown below? Give an exact answer using a square root.



6. What is the volume of the sphere shown below? Give an exact answer using a square root.



Lesson 20: Truncated Cones

Classwork

Opening Exercise 1

1. Examine the bucket below. It has a height of 9 inches and a radius at the top of the bucket of 4 inches.





- a. Describe the shape of the bucket. What is it similar to?

- b. Estimate the volume of the bucket.


Example 1

Determine the volume of the truncated cone shown below.




Exercises 2–6

2. Find the volume of the truncated cone.




- a. Write a proportion that will allow you to determine the height of the cone that has been removed. Explain what all parts of the proportion represent.
- b. Solve your proportion to determine the height of the cone that has been removed.
- c. Write an expression that can be used to determine the volume of the truncated cone. Explain what each part of the expression represents.
- d. Calculate the volume of the truncated cone.

3. Find the volume of the truncated cone.



4. Find the volume of the truncated pyramid with a square base.




- a. Write a proportion that will allow you to determine the height of the cone that has been removed. Explain what all parts of the proportion represent.
- b. Solve your proportion to determine the height of the pyramid that has been removed.
- c. Write an expression that can be used to determine the volume of the truncated pyramid. Explain what each part of the expression represents.
- d. Calculate the volume of the truncated pyramid.

5. A pastry bag is a tool used to decorate cakes and cupcakes. Pastry bags take the form of a truncated cone when filled with icing. What is the volume of a pastry bag with a height of 6 inches, large radius of 2 inches, and small radius of 0.5 inches?
6. Explain in your own words what a truncated cone is and how to determine its volume.

Lesson Summary

A truncated cone or pyramid is a solid figure that is obtained by removing the top portion above a plane parallel to the base. Shown below on the left is a truncated cone. A truncated cone with the top portion still attached is shown below on the right.

Truncated cone:



Truncated cone with top portion attached:




To determine the volume of a truncated cone, you must first determine the height of the portion of the cone that has been removed using ratios that represent the corresponding sides of the right triangles. Next, determine the volume of the portion of the cone that has been removed and the volume of the truncated cone with the top portion attached. Finally, subtract the volume of the cone that represents the portion that has been removed from the complete cone. The difference represents the volume of the truncated cone.

Pictorially,




Problem Set


1. Find the volume of the truncated cone.
 - a. Write a proportion that will allow you to determine the height of the cone that has been removed. Explain what all parts of the proportion represent.
 - b. Solve your proportion to determine the height of the cone that has been removed.
 - c. Show a fact about the volume of the truncated cone using an expression. Explain what each part of the expression represents.
 - d. Calculate the volume of the truncated cone.




2. Find the volume of the truncated cone.



3. Find the volume of the truncated pyramid with a square base.




4. Find the volume of the truncated pyramid with a square base. Note: 3 mm is the distance from the center to the edge of the square at the top of the figure.




5. Find the volume of the truncated pyramid with a square base. Note: 0.5 cm is the distance from the center to the edge of the square at the top of the figure.

6. Explain how to find the volume of a truncated cone.



7. Challenge: Find the volume of the truncated cone.




Lesson 21: Volume of Composite Solids

Classwork


Exercises 1–4

1. a. Write an expression that can be used to find the volume of the chest shown below. Explain what each part of your expression represents.




- b. What is the approximate volume of the chest shown below? Use 3.14 for π . Round your final answer to the tenths place.

2. a. Write an expression that can be used to find the volume of the figure shown below. Explain what each part of your expression represents.




- b. Assuming every part of the cone can be filled with ice cream, what is the exact and approximate volume of the cone and scoop? (Recall that exact answers are left in terms of π and approximate answers use 3.14 for π). Round your approximate answer to the hundredths place.

3. a. Write an expression that can be used to find the volume of the figure shown below. Explain what each part of your expression represents.



- b. Every part of the trophy shown below is made out of silver. How much silver is used to produce one trophy? Give an exact and approximate answer rounded to the hundredths place.

4. Use the diagram of scoops below to answer parts (a) and (b).
- a. Order the scoops from least to greatest in terms of their volumes. Each scoop is measured in inches.




- b. How many of each scoop would be needed to add a half-cup of sugar to a cupcake mixture? (One-half cup is approximately 7 in^3 .) Round your answer to a whole number of scoops.

Lesson Summary


Composite solids are figures that are comprised of more than one solid. Volumes of composite solids can be added as long as no parts of the solids overlap. That is, they touch only at their boundaries.

Problem Set


1. What volume of sand would be required to completely fill up the hourglass shown below? Note: 12m is the height of the truncated cone, not the lateral length of the cone.



2. a. Write an expression that can be used to find the volume of the prism with the pyramid portion removed. Explain what each part of your expression represents.




- b. What is the volume of the prism shown above with the pyramid portion removed?
3. a. Write an expression that can be used to find the volume of the funnel shown below. Explain what each part of your expression represents.




- b. Determine the exact volume of the funnel shown above.

4. What is the approximate volume of the rectangular prism with a cylindrical hole shown below? Use 3.14 for π . Round your answer to the tenths place.



5. A layered cake is being made to celebrate the end of the school year. What is the exact total volume of the cake shown below?




Lesson 22: Average Rate of Change


Classwork

Exercise

The height of a container in the shape of a circular cone is 7.5 ft., and the radius of its base is 3 ft., as shown. What is the total volume of the cone?




Time in minutes	Water level in feet
	1
	2
	3
	4
	5
	6
	7
	7.5



Problem Set


- Complete the table below for more intervals of water levels of the cone discussed in class. Then graph the data on a coordinate plane.

Time in minutes	Water level in feet
	1
	1.5
	2
	2.5
	3
	3.5
	4
	4.5
	5
	5.5
	6
	6.5
	7
	7.5




2. Complete the table below and graph the data on a coordinate plane. Compare the graphs from Problems 1 and 2. What do you notice? If you could write a rule to describe the function of the rate of change of the water level of the cone, what might the rule include?

x	\sqrt{x}
1	
4	
9	
16	
25	
36	
49	
64	
81	
100	



3. Describe, intuitively, the rate of change of the water level if the container being filled were a cylinder. Would we get the same results as with the cone? Why or why not? Sketch a graph of what filling the cylinder might look like, and explain how the graph relates to your answer.




4. Describe, intuitively, the rate of change if the container being filled were a sphere. Would we get the same results as with the cone? Why or why not?

Lesson 23: Nonlinear Motion


Classwork

Exercise





A ladder of length L ft. leaning against a wall is sliding down. The ladder starts off being flush (right up against) with the wall. The top of the ladder slides down the vertical wall at a constant speed of v ft. per second. Let the ladder in the position L_1 slide down to position L_2 after 1 second, as shown below.



Will the bottom of the ladder move at a constant rate away from point O ?



Consider the three right triangles shown below. Specifically the change in the length of the base as the height decreases in increments of 1 ft.



Input t	Output $y = \sqrt{t(30 - t)}$
0	
1	
3	
4	
7	
8	
14	
15	

Problem Set

1. Suppose the ladder is 10 feet long, and the top of the ladder is sliding down the wall at a rate of 0.8 ft. per second. Compute the average rate of change in the position of the bottom of the ladder over the intervals of time from 0 to 0.5 seconds, 3 to 3.5 seconds, 7 to 7.5 seconds, 9.5 to 10 seconds, and 12 to 12.5 seconds. How do you interpret these numbers?

Input t	Output $y = \sqrt{0.8t(20 - 0.8t)}$
0	
0.5	
3	
3.5	
7	
7.5	
9.5	
10	
12	
12.5	

2. Will any length of ladder, L , and any constant speed of sliding of the top of the ladder v ft. per second, ever produce a constant rate of change in the position of the bottom of the ladder? Explain.